

①

HW 5 Solutions
 Math 3150-4

3.7.2 Solve 2D WEQ with $c = \frac{1}{\pi}$ and
 $f(x, y) = \sin \pi x \sin \pi y$ for $0 \leq x \leq 1$ and $0 \leq y \leq 1$
 $g(x, y) = \sin \pi x$

The Fourier series of $f(x, y)$ is itself: $B_{m,n} = \begin{cases} 1 & m=1, n=1 \\ 0 & m \neq 1, n \neq 1 \end{cases}$

This is easy to see since $\int_0^1 \int_0^1 \sin \pi x \sin \pi y \sin m \pi x \sin n \pi y dx dy$ is a 2D inner product

$$B_{m,n} = \frac{(\sin \pi x \sin \pi y, \sin m \pi x \sin n \pi y)}{(\sin \pi x \sin \pi y, \sin \pi x \sin \pi y)} = \frac{(\sin \pi x, \sin m \pi x) (\sin \pi y, \sin n \pi y)}{(\sin \pi x, \sin \pi x) (\sin \pi y, \sin \pi y)}$$

$$= \begin{cases} 1 & \text{if } n=1 \\ 0 & \text{if } n \neq 1 \end{cases} = \begin{cases} 1 & \text{if } m=1 \\ 0 & \text{if } m \neq 1 \end{cases}$$

the Fourier coefficients of $g(x, y)$ are $B_{m,n}^* \sqrt{m^2 + n^2}$

thus:

$$B_{m,n}^* \sqrt{m^2 + n^2} = \frac{(\sin \pi x, \sin m \pi x \sin n \pi y)}{(\sin \pi x \sin \pi y, \sin m \pi x \sin n \pi y)}$$

$$= \frac{(\sin \pi x, \sin m \pi x)}{(\sin \pi x, \sin \pi x)} \frac{(1, \sin n \pi y)}{(\sin \pi y, \sin \pi y)}$$

$$= \begin{cases} 1 & \text{if } n=1 \\ 0 & \text{if } n \neq 1 \end{cases}$$

now $(1, \sin n \pi y) = \int_0^1 1 \sin n \pi y dy = \left. \frac{-\cos n \pi y}{n \pi} \right|_0^1 = \frac{1}{n \pi} (1 - (-1)^n)$

$(\sin \pi y, \sin \pi y) = \int_0^1 \sin^2 \pi y dy = \frac{1}{2} \int_0^1 (1 - \cos 2 \pi y) dy = \frac{1}{2}$









$$\Rightarrow B_{m,n}^* = \begin{cases} \frac{1}{\sqrt{1+m^2}} \frac{1}{2 n \pi} (1 - (-1)^n) & \text{if } n=1 \\ 0 & \text{if } n \neq 1 \end{cases}$$

Putting it all together

$$\psi(x, y, t) = \sin \pi x \sin \pi y \cos(\sqrt{2} t) + \sum_{m=1}^{\infty} \frac{1}{\sqrt{1+m^2}} \frac{1}{2m\pi} (1 - (-1)^m) \sin \pi x \sin m \pi y \sin(\sqrt{1+m^2} t)$$

(note: I had swapped m and n here - but the answer is still correct)

plots:



3.7.13

2D Heat eq. on $[0, 1] \times [0, 1]$ w/
init temp distrib

$$f(x, y) = \sin \pi x \sin \pi y$$

and $c = 1$.The Fourier series of $f(x, y)$ is f itself. $(A_{n,m} = \begin{cases} 1 & \text{if } n=1, m=1 \\ 0 & \text{otherwise} \end{cases})$

$$\Rightarrow u(x, y, t) = \sin \pi x \sin \pi y \exp[-\pi \sqrt{2} t]$$

3.7.16

To prove orthogonality relations in 2D we use these in 1D

$$\text{Let } (u, v) = \int_0^a \int_0^b u(x, y) v(x, y) dx dy$$

then:

$$\begin{aligned} & \left(\sin \frac{m\pi}{a} x \sin \frac{m'\pi}{b} y, \sin \frac{n\pi}{a} x \sin \frac{n'\pi}{b} y \right) \\ &= \left(\sin \frac{m\pi}{a} x, \sin \frac{n\pi}{a} x \right) \left(\sin \frac{m'\pi}{b} y, \sin \frac{n'\pi}{b} y \right) \\ &= \int_0^a \sin \frac{m\pi}{a} x \sin \frac{n\pi}{a} x dx \int_0^b \sin \frac{m'\pi}{b} y \sin \frac{n'\pi}{b} y dy \\ &= \begin{cases} \frac{a}{2} & \text{if } m=n' \\ 0 & \text{otherwise} \end{cases} = \begin{cases} \frac{b}{2} & \text{if } m=n' \\ 0 & \text{otherwise} \end{cases} \end{aligned}$$

$$= \begin{cases} \frac{ab}{4} & \text{if } m=m' \text{ and } n=n' \\ 0 & \text{otherwise} \end{cases}$$

here we used the fact that:

$$\int_0^a \int_0^b f(x) g(y) dy dx = \left(\int_0^a f(x) dx \right) \left(\int_0^b g(y) dy \right)$$

orthogonality relations in 1D.
see e.g. p 22

```

% Math 3150-1
%
% Sample code for problem 3.7.2:
%
% Square membrane with side 1, wave velocity  $c=1/\pi$  and
%   initial shape  $u(x,y,0) = f(x,y) = \sin(\pi x) * \sin(\pi y)$ 
%   initial velocity  $u_t(x,y,0) = g(x,y) = \sin(\pi x)$ 

% number of terms in the expansion
N=30; M=30;

% Time steps and final time
nt=40; Tmax=pi;
ts = linspace(0,Tmax,nt);

% whether to take snapshots for several times
take_snapshots=1;
nsnaps = 8;
snapbasename = 'p3_7_2';
snapcount = 0;
if (mod(nt,nsnaps)~=0)
    error('number_of_snapshots_must_divide_number_of_time_steps');
end;

% setup a grid to plot the function
Nx=20; Ny=20; % number of points in x and y directions
x=linspace(0,1,Nx); y=linspace(0,1,Ny);
[xx,yy]=meshgrid(x,y);

for it=1:length(ts),
    t = ts(it);
    ss = sin(pi*xx).*sin(pi*yy)*cos(sqrt(2)*t);
    for n=1:N,
        lambda1n = sqrt(1+n^2);
        B1nstar = (1-(-1)^n)/(lambda1n*2*n*pi);
        ss = ss + sin(pi*xx).*sin(n*pi*yy) * B1nstar * sin(lambda1n*t);
    end;
    % the rest of the loop handles plotting
    mesh(xx,yy,ss,'edgecolor','k','facecolor','none');
    axis([0 1 0 1 -1 1]);
    title(sprintf('t=%d',t)); xlabel('x'); ylabel('y');
    pause(0.2); % 0.2 is the time we pause
    % take a snapshot if necessary
    if (mod(it-1,nt/nsnaps)==0 & take_snapshots)
        filename = sprintf('%s_%03d.png',snapbasename,snapcount);
        fprintf(['saving_to_file_' filename '\n']);
        print('-dpng','-r50',filename);
        snapcount=snapcount+1;
    end;
end;
end;

```