

```
> # Sample quiz answer checks, maple text only.
```

```
> # answer check problem (a)
```

```
> f:=t*exp(2*t)+2*t*sin(3*t)+3*exp(-t)*cos(4*t);
```

$$f := t e^{2t} + 2 t \sin(3 t) + 3 e^{-t} \cos(4 t) \quad (1)$$

```
> with(inttrans): # load laplace package
```

```
> laplace(f,t,s);
```

$$\frac{1}{(s-2)^2} + \frac{12s}{(s^2+9)^2} + \frac{3}{2(s+1-4I)} + \frac{3}{2(s+1+4I)} \quad (2)$$

```
> # The last two fractions simplify to 3(s+1)/((s+1)^2+16).
```

```
> # answer check problem (b)
```

```
> F:=16/(s^2+4)+(s+1)/(s^2-2*s+10)+2/(s^2+16);
```

$$F := \frac{16}{s^2+4} + \frac{s+1}{s^2-2s+10} + \frac{2}{s^2+16} \quad (3)$$

```
> invlaplace(F,s,t);
```

$$8 \sin(2 t) + \frac{1}{2} \sin(4 t) + \frac{1}{3} e^t (3 \cos(3 t) + 2 \sin(3 t)) \quad (4)$$

```
> # answer check problem (c)
```

```
> de:=diff(x(t),t,t)+256*x(t)=1;ic:=x(0)=1,D(x)(0)=0;
```

$$de := \frac{d^2}{dt^2} x(t) + 256 x(t) = 1$$

$$ic := x(0) = 1, D(x)(0) = 0 \quad (5)$$

```
> dsolve([de,ic],x(t));
```

$$x(t) = \frac{1}{256} + \frac{255}{256} \cos(16 t) \quad (6)$$

```
> # answer check problem (c), partial fractions
```

```
> convert((s^2+1)/(s*(s^2+256)),parfrac,s);
```

$$\frac{1}{256 s} + \frac{255}{256} \frac{s}{s^2+256} \quad (7)$$