

Sample Quiz 8

Sample Quiz 8, Problem 1. Solving Higher Order Constant-Coefficient Equations

The **Algorithm** applies to constant-coefficient homogeneous linear differential equations of order N , for example equations like

$$y'' + 16y = 0, \quad y'''' + 4y'' = 0, \quad \frac{d^5 y}{dx^5} + 2y''' + y'' = 0.$$

1. Find the N th degree characteristic equation by Euler's substitution $y = e^{rx}$. For instance, $y'' + 16y = 0$ has characteristic equation $r^2 + 16 = 0$, a polynomial equation of degree $N = 2$.
2. Find all real roots and all complex conjugate pairs of roots satisfying the characteristic equation. List the N roots according to multiplicity.
3. Construct N distinct Euler solution atoms from the list of roots. Then the general solution of the differential equation is a linear combination of the Euler solution atoms with arbitrary coefficients c_1, c_2, c_3, \dots .

The solution space S of the differential equation is given by

$$S = \text{span}(\text{the } N \text{ Euler solution atoms}).$$

Examples: Constructing Euler Solution Atoms from roots.

Three roots $0, 0, 0$ produce three atoms $e^{0x}, xe^{0x}, x^2e^{0x}$ or $1, x, x^2$.

Three roots $0, 0, 2$ produce three atoms e^{0x}, xe^{0x}, e^{2x} .

Two complex conjugate roots $2 \pm 3i$ produce two atoms $e^{2x} \cos(3x), e^{2x} \sin(3x)$.¹

Four complex conjugate roots listed according to multiplicity as $2 \pm 3i, 2 \pm 3i$ produce four atoms $e^{2x} \cos(3x), e^{2x} \sin(3x), xe^{2x} \cos(3x), xe^{2x} \sin(3x)$.

Seven roots $1, 1, 3, 3, 3, \pm 3i$ produce seven atoms $e^x, xe^x, e^{3x}, xe^{3x}, x^2e^{3x}, \cos(3x), \sin(3x)$.

Two conjugate complex roots $a \pm bi$ ($b > 0$) arising from roots of $(r - a)^2 + b^2 = 0$ produce two atoms $e^{ax} \cos(bx), e^{ax} \sin(bx)$.

The Problem

Solve for the general solution or the particular solution satisfying initial conditions.

(a) $y'' + 16y' = 0$

(b) $y'' + 16y = 0$

(c) $y'''' + 16y'' = 0$

(d) $y'' + 16y = 0, y(0) = 1, y'(0) = -1$

(e) $y'''' + 9y'' = 0, y(0) = y'(0) = 0, y''(0) = y'''(0) = 1$

(f) The characteristic equation is $(r - 2)^2(r^2 - 4) = 0$.

(g) The characteristic equation is $(r - 1)^2(r^2 - 1)((r + 2)^2 + 4) = 0$.

(h) The characteristic equation roots, listed according to multiplicity, are $0, 0, 0, -1, 2, 2, 3 + 4i, 3 - 4i$.

¹The Reason: $\cos(3x) = \frac{1}{2}e^{3xi} + \frac{1}{2}e^{-3xi}$ by Euler's formula $e^{i\theta} = \cos \theta + i \sin \theta$. Then $e^{2x} \cos(3x) = \frac{1}{2}e^{2x+3xi} + \frac{1}{2}e^{2x-3xi}$ is a linear combination of exponentials e^{rx} where r is a root of the characteristic equation. Euler's substitution implies e^{rx} is a solution, so by superposition, so also is $e^{2x} \cos(3x)$. Similar for $e^{2x} \sin(3x)$.

Solutions to Problem 1

(a) $y'' + 16y' = 0$ upon substitution of $y = e^{rx}$ becomes $(r^2 + 16r)e^{rx} = 0$. Cancel e^{rx} to find the **characteristic equation** $r^2 + 16r = 0$. It factors into $r(r + 16) = 0$, then the two roots r make the list $r = 0, -16$. The Euler solution atoms for these roots are e^{0x}, e^{-16x} . Report the general solution $y = c_1 e^{0x} + c_2 e^{-16x} = c_1 + c_2 e^{-16x}$, where symbols c_1, c_2 stand for arbitrary constants.

(b) $y'' + 16y = 0$ has characteristic equation $r^2 + 16 = 0$. Because a quadratic equation $(r - a)^2 + b^2 = 0$ has roots $r = a \pm bi$, then the root list for $r^2 + 16 = 0$ is $0 + 4i, 0 - 4i$, or briefly $\pm 4i$. The Euler solution atoms are $e^{0x} \cos(4x), e^{0x} \sin(4x)$. The general solution is $y = c_1 \cos(4x) + c_2 \sin(4x)$, because $e^{0x} = 1$.

(c) $y'''' + 16y'' = 0$ has characteristic equation $r^4 + 4r^2 = 0$ which factors into $r^2(r^2 + 16) = 0$ having root list $0, 0, 0 \pm 4i$. The Euler solution atoms are $e^{0x}, x e^{0x}, e^{0x} \cos(4x), e^{0x} \sin(4x)$. Then the general solution is $y = c_1 + c_2 x + c_3 \cos(4x) + c_4 \sin(4x)$.

(d) $y'' + 16y = 0, y(0) = 1, y'(0) = -1$ defines a particular solution y . The usual arbitrary constants c_1, c_2 are determined by the initial conditions. From part (b), $y = c_1 \cos(4x) + c_2 \sin(4x)$. Then $y' = -4c_1 \sin(4x) + 4c_2 \cos(4x)$. Initial conditions $y(0) = 1, y'(0) = -1$ imply the equations $c_1 \cos(0) + c_2 \sin(0) = 1, -4c_1 \sin(0) + 4c_2 \cos(0) = -1$. Using $\cos(0) = 1$ and $\sin(0) = 0$ simplifies the equations to $c_1 = 1$ and $4c_2 = -1$. Then the particular solution is $y = c_1 \cos(4x) + c_2 \sin(4x) = \cos(4x) - \frac{1}{4} \sin(4x)$.

(e) $y'''' + 9y'' = 0, y(0) = y'(0) = 0, y''(0) = y'''(0) = 1$ is solved like part (d). First, the characteristic equation $r^4 + 9r^2 = 0$ is factored into $r^2(r^2 + 9) = 0$ to find the root list $0, 0, 0 \pm 3i$. The Euler solution atoms are $e^{0x}, x e^{0x}, e^{0x} \cos(3x), e^{0x} \sin(3x)$, which implies the general solution $y = c_1 + c_2 x + c_3 \cos(3x) + c_4 \sin(3x)$. We have to find the derivatives of y : $y' = c_2 - 3c_3 \sin(3x) + 3c_4 \cos(3x), y'' = -9c_3 \cos(3x) - 9c_4 \sin(3x), y''' = 27c_3 \sin(3x) - 27c_4 \cos(3x)$. The initial conditions give four equations in four unknowns c_1, c_2, c_3, c_4 :

$$\begin{array}{rccccrcr} c_1 & + & c_2(0) & + & c_3 \cos(0) & + & c_4 \sin(0) & = & 0, \\ & & c_2 & - & 3c_3 \sin(0) & + & 3c_4 \cos(0) & = & 0, \\ & & & - & 9c_3 \cos(0) & - & 9c_4 \sin(0) & = & 1, \\ & & & & 27c_3 \sin(0) & - & 27c_4 \cos(0) & = & 1, \end{array}$$

which has invertible coefficient matrix $\begin{pmatrix} 1 & 0 & 1 & 0 \\ 0 & 1 & 0 & 3 \\ 0 & 0 & -9 & 0 \\ 0 & 0 & 0 & -27 \end{pmatrix}$ and right side vector $\begin{pmatrix} 0 \\ 0 \\ 1 \\ 1 \end{pmatrix}$. The

solution is $c_1 = c_2 = 1/9, c_3 = -1/9, c_4 = -1/27$. Then the particular solution is $y = c_1 + c_2 x + c_3 \cos(3x) + c_4 \sin(3x) = \frac{1}{9} + \frac{1}{9}x - \frac{1}{9} \cos(3x) - \frac{1}{27} \sin(3x)$

(f) The characteristic equation is $(r - 2)^2(r^2 - 4) = 0$. Then $(r - 2)^3(r + 2) = 0$ with root list $2, 2, 2, -2$ and Euler atoms $e^{2x}, x e^{2x}, x^2 e^{2x}, e^{-2x}$. The general solution is a linear combination of these four atoms.

(g) The characteristic equation is $(r - 1)^2(r^2 - 1)((r + 2)^2 + 4) = 0$. The root list is $1, 1, 1, -1, -2 \pm 2i$ with Euler atoms $e^x, x e^x, x^2 e^x, e^{-x}, e^{-2x} \cos(2x), e^{-2x} \sin(2x)$. The general solution is a linear combination of these six atoms.

(h) The characteristic equation roots, listed according to multiplicity, are $0, 0, 0, -1, 2, 2, 3 + 4i, 3 - 4i$. Then the Euler solution atoms are $e^{0x}, x e^{0x}, x^2 e^{0x}, e^{-x}, e^{2x}, x e^{2x}, e^{3x} \cos(4x), e^{3x} \sin(4x)$. The general solution is a linear combination of these eight atoms.

Sample Quiz 8, Problem 2. Laplace Theory

Laplace theory implements the *method of quadrature* for higher order differential equations, linear systems of differential equations, and certain partial differential equations.

Laplace's method solves **differential equations**.

The Problem. Solve by table methods or Laplace's method.

(a) Forward table. Find $\mathcal{L}(f(t))$ for $f(t) = te^{2t} + 2t \sin(3t) + 3e^{-t} \cos(4t)$.

(b) Backward table. Find $f(t)$ for

$$\mathcal{L}(f(t)) = \frac{16}{s^2 + 4} + \frac{s + 1}{s^2 - 2s + 10} + \frac{2}{s^2 + 16}.$$

(c) Solve the initial value problem $x''(t) + 256x(t) = 1$, $x(0) = 1$, $x'(0) = 0$.

Solution (a).

$$\begin{aligned} \mathcal{L}(f(t)) &= \mathcal{L}(te^{2t} + 2t \sin(3t) + 3e^{-t} \cos(4t)) \\ &= \mathcal{L}(te^{2t}) + 2\mathcal{L}(t \sin(3t)) + 3\mathcal{L}(e^{-t} \cos(4t)) && \text{Linearity} \\ &= -\frac{d}{ds}\mathcal{L}(e^{2t}) - 2\frac{d}{ds}\mathcal{L}(\sin(3t)) + 3\mathcal{L}(e^{-t} \cos(4t)) && \text{Differentiation rule} \\ &= -\frac{d}{ds}\mathcal{L}(e^{2t}) - 2\frac{d}{ds}\mathcal{L}(\sin(3t)) + 3\mathcal{L}(\cos(4t))\Big|_{s=s+1} && \text{Shift rule} \\ &= -\frac{d}{ds}\frac{1}{s-2} - 2\frac{d}{ds}\frac{3}{s^2+9} + 3\frac{s}{s^2+16}\Big|_{s=s+1} && \text{Forward table} \\ &= \frac{1}{(s-2)^2} + \frac{12s}{(s^2+9)^2} + 3\frac{s+1}{(s+1)^2+16} && \text{Calculus} \end{aligned}$$

Solution (b).

$$\begin{aligned} \mathcal{L}(f(t)) &= \frac{16}{s^2+4} + \frac{s+1}{s^2-2s+10} + \frac{2}{s^2+16} \\ &= 8\frac{2}{s^2+4} + \frac{s+1}{(s-1)^2+9} + \frac{1}{2}\frac{4}{s^2+16} && \text{Prep for backward table} \\ &= 8\mathcal{L}(\sin 2t) + \frac{s+1}{(s-1)^2+9} + \frac{1}{2}\mathcal{L}(\sin 4t) && \text{backward table} \\ &= 8\mathcal{L}(\sin 2t) + \frac{s+2}{s^2+9}\Big|_{s=s-1} + \frac{1}{2}\mathcal{L}(\sin 4t) && \text{shift rule} \\ &= 8\mathcal{L}(\sin 2t) + \mathcal{L}(\cos 3t + \frac{2}{3}\sin 3t)\Big|_{s=s-1} + \frac{1}{2}\mathcal{L}(\sin 4t) && \text{backward table} \\ &= 8\mathcal{L}(\sin 2t) + \mathcal{L}(e^t \cos 3t + e^t \frac{2}{3}\sin 3t) + \frac{1}{2}\mathcal{L}(\sin 4t) && \text{shift rule} \\ &= \mathcal{L}(8 \sin 2t) + e^t \cos 3t + e^t \frac{2}{3}\sin 3t + \frac{1}{2}\mathcal{L}(\sin 4t) && \text{Linearity} \\ f(t) &= 8 \sin 2t + e^t \cos 3t + e^t \frac{2}{3}\sin 3t + \frac{1}{2}\sin 4t && \text{Lerch's cancel rule} \end{aligned}$$

Solution (c).

$$\begin{aligned} \mathcal{L}(x''(t) + 256x(t)) &= \mathcal{L}(1) && \mathcal{L} \text{ acts like matrix mult} \\ s\mathcal{L}(x') - x'(0) + 256\mathcal{L}(x) &= \mathcal{L}(1) && \text{Parts rule} \\ s(s\mathcal{L}(x) - x(0)) - x'(0) + 256\mathcal{L}(x) &= \mathcal{L}(1) && \text{Parts rule} \\ s^2\mathcal{L}(x) - s + 256\mathcal{L}(x) &= \mathcal{L}(1) && \text{Use } x(0) = 1, x'(0) = 0 \\ (s^2 + 256)\mathcal{L}(x) &= s + \mathcal{L}(1) && \text{Collect } \mathcal{L}(x) \text{ left} \end{aligned}$$

$$\begin{aligned} \mathcal{L}(x) &= \frac{s+\mathcal{L}(1)}{(s^2+256)} && \text{Isolate } \mathcal{L}(x) \text{ left} \\ \mathcal{L}(x) &= \frac{s+1/s}{(s^2+256)} && \text{Forward table} \\ \mathcal{L}(x) &= \frac{s^2+1}{s(s^2+256)} && \text{Algebra} \\ \mathcal{L}(x) &= \frac{A}{s} + \frac{Bs+C}{s^2+256} && \text{Partial fractions} \\ \mathcal{L}(x) &= A\mathcal{L}(1) + B\mathcal{L}(\cos 16t) + \frac{C}{16}\mathcal{L}(\sin 16t) && \text{Backward table} \\ \mathcal{L}(x) &= \mathcal{L}(A + B \cos 16t + \frac{C}{16} \sin 16t) && \text{Linearity} \\ x(t) &= A + B \cos 16t + \frac{C}{16} \sin 16t && \text{Lerch's rule} \end{aligned}$$

The partial fraction problem remains:

$$\frac{s^2 + 1}{s(s^2 + 256)} = \frac{A}{s} + \frac{Bs + C}{s^2 + 256}$$

This problem is solved by clearing the fractions, then swapping sides of the equation, to obtain

$$A(s^2 + 256) + (Bs + C)(s) = s^2 + 1.$$

Substitute three values for s to find 3 equations in 3 unknowns A, B, C :

$$\begin{aligned} s = 0 & \quad 256A & = & 1 \\ s = 1 & \quad 257A + B + C & = & 2 \\ s = -1 & \quad 257A + B - C & = & 2 \end{aligned}$$

Then $A = 1/256, B = 255/256, C = 0$ and finally

$$x(t) = A + B \cos 16t + \frac{C}{16} \sin 16t = \frac{1 + 255 \cos 16t}{256}$$

Answer Checks

```
# Sample quiz 8
# answer check problem 2(a)
f:=t*exp(2*t)+2*t*sin(3*t)+3*exp(-t)*cos(4*t);
with(inttrans): # load laplace package
laplace(f,t,s);
# The last two fractions simplify to 3(s+1)/((s+1)^2+16).
# answer check problem 2(b)
F:=16/(s^2+4)+(s+1)/(s^2-2*s+10)+2/(s^2+16);
invlaplace(F,s,t);
# answer check problem 2(c)
de:=diff(x(t),t,t)+256*x(t)=1;ic:=x(0)=1,D(x)(0)=0;
dsolve([de,ic],x(t));
# answer check problem 2(c), partial fractions
convert((s^2+1)/(s*(s^2+256)),parfrac,s);
```

The output appears on the next page

```

[> # Sample quiz 11
[> # answer check problem 2(a)
> f:=t*exp(2*t)+2*t*sin(3*t)+3*exp(-t)*cos(4*t);
      f:= t e2t + 2 t sin(3 t) + 3 e-t cos(4 t) (1)
[> with(inttrans): # load laplace package
> laplace(f,t,s) assuming s::real;
      1      12 s      3      3
      (s-2)2 + (s2+9)2 + 2(s+1-4I) + 2(s+1+4I) (2)
[> # The last two fractions simplify to 3(s+1)/((s+1)^2+16).
[> # answer check problem 2(b)
> F:=16/(s^2+4)+(s+1)/(s^2-2*s+10)+2/(s^2+16);
      F:= 16      s+1      2
          s2+4 + s2-2s+10 + s2+16 (3)
[> invlaplace(F,s,t);
      8 sin(2 t) + 1/2 sin(4 t) + 1/3 et (3 cos(3 t) + 2 sin(3 t)) (4)
[> # answer check problem 2(c)
> de:=diff(x(t),t,t)+256*x(t)=1;ic:=x(0)=1,D(x)(0)=0;
      de:= d2 x(t) + 256 x(t) = 1
          dt2
      ic:= x(0) = 1, D(x)(0) = 0 (5)
[> dsolve([de,ic],x(t));
      x(t) = 1/256 + 255/256 cos(16 t) (6)
[> # answer check problem 2(c), partial fractions
> convert((s^2+1)/(s*(s^2+256)),parfrac,s);
      1      255      s
      256 s + 256 s2+256 (7)

```