

Math 2250-3

Wed 9/29

Determinants § 3.6

HW for Monday 10/4

§ 3.6 2, 6, 17 21, 22

this material will be on the midterm

In 22 & 32

1

just find x_1 with Cramer's rule. Then compute A^{-1} with adjoint formula (page 210), as well as with the rref algorithm (better get same ans!)

well as with the rref algorithm (better get same ans!)

Finally, use A^{-1} to solve $A\vec{x} = \vec{b}$ and compare to Cramer's rule value for x_1

The determinant of a square matrix is a magic number, obtained by strange rules

The matrix is invertible exactly when $\det(A) \neq 0$

exactly when $\text{rref}(A) = I$

When $\det A \neq 0$ there is an algebraic formula for A^{-1} , (adjoint formula) which results in a determinant formula (Cramer's rule) for finding solutions to $A\vec{x} = \vec{b}$

Illustrate for 2×2 matrices:

(then spend today and at least part of Friday discussing $n \times n$ case!)

$$\begin{vmatrix} a & b \\ c & d \end{vmatrix} := ad - bc$$

bars rather than brackets mean take determinant.

$$\text{if } \det A \neq 0 \text{ then } A^{-1} = \frac{1}{\det(A)} \begin{bmatrix} d & -b \\ -c & a \end{bmatrix} \quad \text{check!}$$

example: solve

$$\begin{aligned} 3x + 4y &= 4 \\ 5x + 6y &= -2 \end{aligned}$$

determinant can be defined inductively (i.e. for matrices $(n+1) \times (n+1)$ if you know $n \times n$ det's)

3x3 case:

$$\begin{vmatrix} a_{11} & a_{12} & a_{13} \\ a_{21} & a_{22} & a_{23} \\ a_{31} & a_{32} & a_{33} \end{vmatrix} = a_{11} \begin{vmatrix} a_{22} & a_{23} \\ a_{32} & a_{33} \end{vmatrix} - a_{12} \begin{vmatrix} a_{21} & a_{23} \\ a_{31} & a_{33} \end{vmatrix} + a_{13} \begin{vmatrix} a_{21} & a_{22} \\ a_{31} & a_{32} \end{vmatrix}$$

\uparrow M_{11} (1-1 minor) cross out row₁ & col₁, take 2x2 det
 \uparrow M_{12}
 \uparrow M_{13}


example:

$$\begin{vmatrix} 1 & 2 & -1 \\ 0 & 3 & 1 \\ 2 & -2 & 1 \end{vmatrix}$$

=

$$= a_{11}(a_{22}a_{33} - a_{32}a_{23}) - a_{12}(a_{21}a_{33} - a_{31}a_{23}) + a_{13}(a_{21}a_{32} - a_{31}a_{22})$$

$$= a_{11}a_{22}a_{33} + a_{12}a_{23}a_{31} + a_{13}a_{21}a_{32} - a_{31}a_{22}a_{13} - a_{32}a_{23}a_{11} - a_{33}a_{21}a_{12}$$



"cross hatch" only works for 3x3!

in fact, using matrix of $(-1)^{i+j} = \begin{bmatrix} + & - & + \\ - & + & - \\ + & - & + \end{bmatrix}$

you can compute a 3x3 det by expanding across any row or down any column

$$\det A = \sum_{j=1}^3 (-1)^{i+j} a_{ij} M_{ij} = \sum_{j=1}^3 a_{ij} C_{ij} \quad [i \text{ fixed}]$$

expansion across row i

$$= \sum_{i=1}^3 (-1)^{i+j} a_{ij} M_{ij} = \sum_{i=1}^3 a_{ij} C_{ij} \quad [j \text{ fixed}]$$

down column j

recompute $\begin{vmatrix} 1 & 2 & -1 \\ 0 & 3 & 1 \\ 2 & -2 & 1 \end{vmatrix}$

by expanding down column 1
and by cross-hatching.
Get same answer!

Inductive def'n of det A:
(recursive)

$$\det A := \sum_{j=1}^n (-1)^{1+j} a_{1j} M_{1j} = \sum_{j=1}^n (-1)^{1+j} a_{1j} C_{1j}$$

(expansion across top row)

~~Matrix~~

Theorem (true, but not easy, see appendix for details)

You can compute det A by expanding across
any row or down any column:

$$|A| = \sum_{j=1}^n a_{ij} C_{ij} \quad (\text{fixed } i)$$

$$= \sum_{i=1}^n a_{ij} C_{ij} \quad (\text{fixed } j)$$

("ij Minor")
 $M_{ij} := (n-1) \times (n-1)$ det
 of matrix obtained
 from A by deleting
 $\text{row } i(A)$ & $\text{col } j(A)$
 $C_{ij} := (-1)^{i+j} M_{ij}$
 ("ij cofactor")

Example

$$\begin{vmatrix} 1 & 38 & 106 & 3 \\ 0 & 2 & 92 & 2 \\ 0 & 0 & 3 & 1 \\ 0 & 0 & 0 & 4 \end{vmatrix}$$

$$\begin{vmatrix} 1 & 0 & 0 & 0 \\ 0 & 2 & 0 & 0 \\ 96 & \pi & 3 & 0 \\ 1221 & 34 & 17 & 4 \end{vmatrix}$$

Theorem: If A is upper or lower triangular
then $|A|$ is just the product of the diagonal
entries

Computational Shortcuts:

(and important for understanding too!)

Effects of elementary row ops on determinants:
(analogous results hold for elementary column operations)

(1) swapping two rows changes the sign of the determinant

proof: $\begin{vmatrix} a & b \\ c & d \end{vmatrix} = ad - bc$; $\begin{vmatrix} c & d \\ a & b \end{vmatrix} = cb - ad$

so true for 2x2.

Now it is easy to check general case by induction

n=3 → expand across the row that wasn't swapped and use the n=2 result on each minor.

In general, for $A_{(n+1) \times (n+1)}$ expand across an unswapped row and use the (inductive) assumption that result is true for nxn matrices.

(1b) So, if 2 rows are equal, det = 0

proof: let $\det(A) = x$
if we swap rows the new det is $-x$
but rows were the same, so had to get same det,
i.e. $-x = x \Rightarrow x = 0$.

(2) multiplying a single row by c, multiplies the det by c.

proof: if you multiplied row_i(A) by c, expand new matrix det across row_i:

$$\begin{vmatrix} R_1 \\ R_2 \\ cR_i \\ \vdots \\ R_n \end{vmatrix} = \sum_{j=1}^n (c a_{ij}) C_{ij} = c \sum_{j=1}^n a_{ij} C_{ij} = c \det A.$$

(3) Coolest property: replacing row_i(A) with row_i(A) + c row_k(A) Does Not change det!

proof: we'll expand across row_i(A):

$$\begin{aligned} \text{row}_i \rightarrow \begin{vmatrix} R_1 \\ R_2 \\ R_i + cR_k \\ \vdots \\ R_n \end{vmatrix} &= \sum_{j=1}^n (a_{ij} + c a_{kj}) C_{ij} \\ &= \sum_{j=1}^n a_{ij} C_{ij} + c \sum_{j=1}^n a_{kj} C_{ij} \\ &= \det(A) + c \begin{vmatrix} R_1 \\ R_2 \\ R_k \\ \vdots \\ R_k \\ \vdots \\ R_n \end{vmatrix} \end{aligned}$$

→ 0 by (1b)
← kth row
← ith row

example

$$\begin{vmatrix} 1 & 2 & -1 \\ 0 & 3 & 1 \\ 2 & -2 & 1 \end{vmatrix} \neq$$

$$= \begin{vmatrix} 1 & 2 & -1 \\ 0 & 3 & 1 \\ 0 & -6 & 3 \end{vmatrix} \quad -2R_1 + R_3$$

$$= \begin{vmatrix} 1 & 2 & -1 \\ 0 & 3 & 1 \\ 0 & 0 & 5 \end{vmatrix} \quad 2R_2 + R_3$$

$$= 15 !$$

Corollary of idea (Fri-)

$\det A \neq 0$ iff $\text{rref}(A) = I$ iff A^{-1} exists.